Supplementary data

Table 2. Hazard rate ratios (HRR) and 5-year revision rate estimates, for revision due to deep infection, with number of primary THAs included and number of revisions due to deep infection reported, for the 3 time periods and the 4 fixation methods

| THA fixation Period | No. of THAs | Number revised | | p-value | Number revised ^a | 5-year follow-up Adjusted revision rate (CI) | At risk after 5 years |
|------------------------|----------------|-------------------|---------------|---------|--------------------------------|--|--------------------------|
| All | | | | | | | |
| 2005-2009 | 30,668 | 348 | 1 | | 279 | 0.8 (0.7-0.9) | 29,740 |
| 2010-2014 | 35,338 | 487 | 1.4 (1.2-1.6) | < 0.001 | 465 | 1.2 (1.1–1.3) | 34,130 |
| 2015-2019 | 42,848 | 530 | 1.6 (1.3–1.9) | < 0.001 | 530 | n.a.` | 32,334 |
| Cemented | | | , | | | | |
| 2005-2009 | 20,031 | 237 | 1 | | 190 | 0.9 (0.8-1.1) | 19,389 |
| 2010-2014 | 11,478 | 188 | 1.4 (1.2-1.7) | 0.001 | 180 | 1.4 (1.2–1.6) | 11,022 |
| 2015-2019 | 11,263 | 144 | 1.5 (1.1–1.9) | 0.002 | 144 | n.a. | 8,534 |
| Uncemented | | | | | | | |
| 2005-2009 | 5,008 | 63 | 1 | | 48 | 0.8 (0.5-1.0) | 4,858 |
| 2010-2014 | 9,495 | 121 | 1.2 (0.7-1.7) | 0.3 | 116 | 1.1 (0.9–1.3) | 9,192 |
| 2015-2019 | 16,530 | 392 | 1.6 (1.2-2.3) | 0.005 | 372 | n.a. | 11,964 |
| Reverse hybrid | | | | | | | |
| 2005–2009 | 5,026 | 44 | 1 | | 38 | 0.7 (0.5–1.0) | 4,909 |
| 2010–2014 | 13,660 | 165 | 1.3 (0.9–1.9) | 0.1 | 156 | 0.9 (0.8–1.1) | 13,244 |
| 2015–2019 | 11,464 | 125 | 1.5 (1.0–2.2) | 0.04 | 125 | n.a. | 9,481 |
| Hybrid | | | | | | | |
| 2005–2009 | 603 | 4 | 1 | | 3 | 0.3 (0.0–3.1) | 584 |
| 2010–2014 | 705 | 13 | 3.6 (1.1–12) | 0.04 | 13 | 1.4 (0.1–13) | 672 |
| 2015–2019 | 3,591 | 53 | 3.6 (1.1–12) | 0.04 | 53 | n.a. | 2,355 |
| | | | | | | | |

^a Due to infection

The HRR and revision estimates are adjusted for sex, age, ASA class, indication for primary THA, duration of surgery, surgical approach, and modularity of the THA. HRRs are given with 95% confidence intervals (CI).

n.a. = not applicable due to incomplete 5-year follow up.

Table 3. HRR with 95% confidence intervals for revision due to infection relative to timespan postoperatively, for the 3 time periods

| Period | THAs overall | Revised ^a overall | | -30 days d ^a HRR (CI) | | -90 days d ^a HRR (CI) | ays-1 year d ^a HRR (CI) | | -5 years d ^a HRR (CI) | THAs at risk after 5 years |
|--|----------------------------|------------------------------|-------------------|-------------------------------------|----------------|-------------------------------------|---------------------------------------|----------------|-------------------------------------|-------------------------------|
| 2005–2009 2010–2014 2015–2019 ^b | 30,668 35,338 42,848 | 348 487 530 | 103 283 327 | 1 2.2 (1.8–2.8) 2.3 (1.8–2.9) | 41 54 83 | 1 1.0 (0.7–1.6) 1.6 (1.0–2.5) | 1 1.1 (0.7–1.8) 1.6 (1.0–2.6) | 94 76 48 | 1 0.7 (0.5–1.0) 0.9 (0.6–1.3) | , |

^a Due to infection.

The HRR and revision estimates are adjusted for sex, age, ASA class, indication for primary THA, duration of surgery, surgical approach, and modularity of the THA. HRRs are given with 95% confidence intervals.

^b The number revised due to infection will be an underestimate due to incomplete 5-year follow-up.

Table 4. Distribution of patient and surgery related factors, in the 3 periods

| | No. of | Dietribut | ion of fac | tors (%) | |
|----------------------------------|-----------------|---|------------|----------|--|
| | THAs | Distribution of factors (%) 2005– 2010– 2015– | | | |
| Risk factor | 111/13 | 2009 | 2014 | 2019 | |
| 1 lisk lactor | | 2000 | 2014 | 2010 | |
| Total number | 108,854 | 30,668 | 35,338 | 42,848 | |
| Sex | , | , | , | , | |
| Male | 37,710 | 33 | 35 | 36 | |
| Female | 71,144 | 67 | 65 | 64 | |
| Age | · | | | | |
| < 45 | 3,490 | 3 | 3 | 3 | |
| 45–54 | 8,909 | 7 | 8 | 9 | |
| 55–64 | 24,981 | 23 | 23 | 22 | |
| 65–74 | 39,144 | 33 | 36 | 38 | |
| 75–84 | 27,258 | 28 | 25 | 23 | |
| ≥ 85 | 5,072 | 5 | 5 | 5 | |
| ASA class | | | | | |
| 1 | 20,682 | 29 | 16 | 14 | |
| 2 | 66,407 | 52 | 64 | 65 | |
| 3 | 21,318 | 19 | 19 | 20 | |
| 4 | 447 | 0.4 | 0.3 | 0.5 | |
| Indication for primary Th | ΗA | | | | |
| Osteoarthritis | 83,770 | 76 | 77 | 77 | |
| Inflammatory | | | | | |
| hip disease | 2,346 | 3 | 2 | 2 | |
| Acute hip fracture | 3,769 | 2 | 3 | 5 | |
| Complications after | | _ | _ | | |
| hip fracture | 5,743 | 7 | 5 | 4 | |
| childhood hip diseas | | 9 | 9 | 8 | |
| Avascular necrosis of | | 0 | 0 | 0 | |
| the femoral head | 2,888 | 3 | 3 | 3 | |
| Other | 714 | 1 | 1 | 1 | |
| Duration of surgery < 70 minutes | 200 700 | 18 | 27 | 34 | |
| 70–99 minutes | 29,709 | 44 | 42 | 34 41 | |
| 100–129 minutes | 46,287 | 27 | 22 | 18 | |
| ≥ 130 minutes | 24,126 8,732 | 10 | 8 | 6 | |
| Surgical approach | 0,732 | 10 | 0 | O | |
| Anterior | 5,979 | 1 | 6 | 8 | |
| Anterolateral | 8,247 | 6 | 12 | 14 | |
| Lateral | 47,048 | 67 | 50 | 12 | |
| Posterolateral | 47,580 | 25 | 32 | 66 | |
| Modularity of THA | 47,000 | 20 | 02 | 00 | |
| Monobloc | 3,936 | 11 | 2 | 0 | |
| Modular | 104,918 | 89 | 98 | 100 | |
| Fixation principle | 10 1,0 10 | 00 | 00 | 100 | |
| Cemented | 42,772 | 65 | 32 | 26 | |
| Uncemented | 31,033 | 16 | 27 | 39 | |
| Reverse hybrid | 30,150 | 16 | 39 | 27 | |
| Hybrid | 4,899 | 2 | 2 | 8 | |
| | | | | | |

Table 5. Number of primary THAs included and number of reported first revisions due to deep infection, and the association between potential risk factors and revision due to infection

| Risk factor | No. of THAs | Number revised ^a | HRR (CI) | Adjusted 5-year revision rate ^a (CI) |
|---------------------------|----------------|--------------------------------|---------------|---|
| Total number | 108,854 | 1,365 | | |
| Sex | | | | |
| Male | 37,710 | 710 | 2.1 (1.9-2.3) | 1.7 (1.6-1.9) |
| Female | 71,144 | 655 | 1 ` ′ | 0.8 (0.8-0.9) |
| Age | , | | | (/ |
| < 45 | 3,490 | 45 | 1.1 (0.8-1.6) | 1.1 (0.7–1.5) |
| 45–54 | 8,909 | 86 | 0.9 (0.7–1.1) | 0.9 (0.7–1.1) |
| 55-64 | 24,981 | 289 | 1.0 (0.9–1.2) | 1.0 (0.9–1.2) |
| 65–74 | 39,144 | 452 | 1 | 1.0 (0.9–1.1) |
| 75–84 | 27,258 | 397 | 1.2 (1.1-1.4) | 1.2 (1.1–1.4) |
| ≥ 85 | 5,072 | 96 | 1.5 (1.2–1.9) | 1.5 (1.2–1.8) |
| ASA class | - , - | | - (/ | - (/ |
| 1 | 20,682 | 160 | 1 | 0.7 (0.6-0.8) |
| 2 | 66,407 | 770 | 1.5 (1.3–1.8) | 1.1 (1.0–1.2) |
| 3 | 21,318 | 422 | 2.3 (1.9–2.8) | 1.6 (1.4–1.8) |
| 4 | 447 | 13 | 3.3 (1.8-5.8) | |
| Indication for primary TI | | | () | |
| Osteoarthritis | 83,770 | 1,019 | 1 | 1.1 (1.0-1.2) |
| Inflammatory hip | , - | , | | (- / |
| disease | 2,346 | 38 | 1.3 (0.9-1.8) | 1.3 (0.9-1.7) |
| Acute hip fracture | 3,769 | 53 | 1.1 (0.8–1.4) | 1.1 (0.8–1.4) |
| Complications after | -, | | (/ | (/ |
| hip fracture | 5,743 | 104 | 1.2 (1.0-1.5) | 1.3 (1.0-1.5) |
| childhood hip | • | | ` , | ` , |
| disease | 9,624 | 78 | 0.7 (0.6-0.9) | 0.8 (0.6-1.0) |
| Avascular necrosis | , | | , , | , |
| of the femoral head | 2,888 | 60 | 1.5 (1.1-1.9) | 1.8 (1.3-2.3) |
| Other | 714 | 13 | 1.1 (0.6–2.0) | 1.7 (0.6–2.7) |
| Duration of surgery | | | , | , |
| < 70 minutes | 29,709 | 330 | 1.0 (0.8-1.1) | 1.0 (0.9-1.1) |
| 70-99 minutes | 46,287 | 547 | 1 | 1.0 (0.9–1.1) |
| 100-129 minutes | 24,126 | 329 | 1.1 (1.0-1.3) | 1.2 (1.0-1.3) |
| ≥130 minutes | 8,732 | 159 | 1.4 (1.2–1.7) | 1.4 (1.2–1.7) |
| Surgical approach | | | | |
| Anterior | 5,979 | 52 | 0.8 (0.6-1.1) | 0.7 0.5-1.0) |
| Anterolateral | 8,247 | 81 | 1.0 (0.8-1.2) | 0.9 (0.7-1.1) |
| Lateral | 47,048 | 683 | 1.4 (1.2-1.6) | 1.4 (1.3-1.5) |
| Posterolateral | 47,580 | 549 | 1 | 0.9 (0.8-1.0) |
| Modularity of THA | | | | |
| Monobloc | 3,936 | 28 | 0.5 (0.3-0.7) | 0.6 (0.4-0.8) |
| Modular | 104,918 | 1,337 | 1 ` ′ | 1.1 (1.0–1.2) |
| Fixation principle | | | | |
| Cemented | 42,772 | 569 | 1 | 1.0 (0.9-1.2) |
| Uncemented | 31,033 | 392 | 1.2 (1.0-1.4) | 1.3 (1.1–1.4) |
| Reverse hybrid | 30,150 | 334 | 0.8 (0.7-1.0) | 0.9 (0.8–1.0) |
| Hybrid | 4,899 | 70 | 1.3 (1.0–1.6) | 1.3 (1.0–1.7) |

Adjusted revision rates and HRR estimates for sex, age, ASA class, indication for primary THA, duration of surgery, surgical approach, modularity of the THA, and fixation method. The HRRs are presented with 95% confidence intervals. The risk factors are adjusted for all the other risk factors in addition to year of primary surgery to assess the association with revision due to infection.